

IN THE SPECIFICATION:

Please replace the paragraph beginning on page 2, line 23 with the following:

The rabbit skin of the present invention possesses 0.5 iu/g SART activity or more, and which also possesses the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 4, line 17 with the following:

The ~~kallikrein-protease~~kallikrein production inhibition activity referred in this description is determined as follows:

Please replace the paragraph beginning on page 4, line 19 with the following:

The rabbit skin was cut in pieces of 1 cm², 4 times (w/w) of 3% phenol aqueous solution was added, then placed the mixture was placed at 4°C for 72h, and centrifuged after liquid changed into an emulsion. The supernatant was filtrated to collect brown solution A. The brown solution A was boiled for 40 min in a water bath after pH was adjusted to 5.0 by 1M HCl, cooled to 28°C promptly, centrifuged and filtrated to collect solution B. The solution B was boiled for 40 min in a water bath after pH of filtrate was adjusted to 9.2 by 1M NaOH, cooled to 28°C promptly, and filtrated to collect solution C. The pH of the

solution C was adjusted to 4.5 by 1M HCl, activated charcoal was added at 30°C, stirred continuously for 4h, ~~stopped~~ After stirring was terminated, the solution was left it for 30min, ~~removed the~~ The supernatant; was then removed and filtrated under a nitrogen atmosphere. Then, the activated charcoal was dipped in injectable water and washed, filtrated and ~~disarded~~ the filtrate was discarded to collect the activated charcoal, ~~reserved~~ while reserving the activated charcoal. The activated charcoal was then put into injectable water, adjusted pH was adjusted to 11.0 by 1M NaOH, the solution was stirred continuously for 4h, filtrated with a 0.45-µm Millipore filter under nitrogen atmosphere, ~~and washed~~ the activated charcoal was washed by injectable water to collect solution D. After pH of the solution D was adjusted to 6.0 by 1M HCl, the vessel was sealed, heated up and ~~kept the~~ temperature maintained at 121°C for 20min, and then cooled down to under 40°C to collect solution E. The solution E was placed into a decompression distillator, ~~replaced the air was~~ replaced with nitrogen in the decompression distillator, the solution was distilled at 60°C under decompression, and filtrated to collect a solution containing biologically active substances, whose SART activity was determined. The SART activity of the solution was adjusted to 1.2iu/ml by evaporating, concentrating and diluting with distilled water. Next, 10ml of this solution was desalted at 10 µs/cm of final conductance, and dried under decompression condition, into which 1.5 ml of 0.25 M NaCl solution was added to obtain a test solution. Next, 0.2 ml of 0.25M NaCl solution was regarded as the control solution and

treated with a parallel process with the 0.2 ml test solution. Then, 0.5 ml of human plasma ~~were~~was added respectively to both solutions, placed at the freezing point for 5 min, ~~added~~ 0.25 ml of suspension of argilla was added, placed at the freezing point for 20 min, and filtrated. 0.1 ml of filtrate was mixed with 0.2 ml of 0.1 M Tris-HCl buffer and 0.1 ml of basic solution. ~~The~~Effect~~ed~~ reaction was effected for 20 min at 30°C, and was stopped ~~the~~ reaction by adding 0.8 ml of 1% citric acid. The absorbance A of the test solution was determined in a control solution with 405 nm as the absorbance ~~of control solution and~~ was initialized as 0.4. If A was less than 0.4, the rabbit skin from which the test solution was prepared was regarded as possessing the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 6, line 15 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Then, 5 ml of the virus solution was added ~~into~~to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^9 virus/ml. A Japanese white rabbit, weighing 3 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.4 ml per site in 200 sites with no leaking, no injecting without the virus injection and no puncturing throughout the skin. The injected rabbit ~~injected~~ was fed for 4 days. When the inflammatory

tissue showed that the skin surface had visible blains ~~accompanying with~~ accompanied by changing colour from redness to mauveness, ~~and the~~ skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra dislocation and it was peeled ~~inafter~~ 15 min. The skin of the rabbit was packed in a plastic bag, and was preserved at -18°C in a refrigerator prior to use. The rabbit skin weighed 349 g, its SART activity was 0.85 iu/g, and its absorbance was 0.07. The results indicated that it possessed the kallikrein-protease kallikrein production inhibition activity.

Please replace the paragraph beginning on page 7, line 7 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~ to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^9 virus/ml. A New Zealand white rabbit, weighing 2.75 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.3 ml per site in 250 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying with~~ accompanied by changing colour from redness to mauveness, ~~and the~~ skin became thick, and the subcuticle and hip became swollen, ~~killed the rabbit~~ was killed by dislocating the

cervical vertebra dislocation and it was peeled in after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 302-g, its SART activity was 0.60 iu/g, and its absorbance was 0.1. The results indicated that it possessed the ~~kallikrein-protease~~ kallikrein production inhibition activity.

Please replace the paragraph beginning on page 7, line 24 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~ to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^6 virus/ml. A Chinese rabbit weighing 1.5 kg, with its back shaved, was sterilized with 75 % ethanol, injected subcutaneously with the virus injection of 0.1 ml per site in 250 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying with~~ accompanied by changing colour from redness to mauveness, ~~and~~ the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by cervical vertebra dislocation and it was peeled ~~in~~ after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 176 g, its

SART activity was 0.50 iu/g, and its absorbance was 0.15. The results indicated that it possessed the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 8, line 15 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^7 virus/ml. A Blue-violet rabbit, weighing 2 kg with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.2 ml per site in 100 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When inflammatory tissue showed that the skin surface had visible blains ~~accompanying with~~accompanied by changing colour from redness to mauveness, ~~and~~the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra ~~dislocation~~ and it was peeled ~~in~~after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 230 g, its SART activity was 0.55 iu/g, and its absorbance was 0.12. The results indicated that it possessed the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 9, line 6 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^9 virus/ml. A New Zealand white rabbit, weighing 2.75 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.3 ml per site in 200 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying~~ with accompanied by changing colour from redness to mauveness, ~~and the~~ skin became thick, and the subcuticle and the hip became swollen, the rabbit was killed by dislocating the cervical vertebra ~~dislocation~~ and it was peeled ~~in~~ after 15 min. The skin of rabbit was packed in a plastic bag, and it was preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 310 g, its SART activity was 0.79 iu/g, and its absorbance was 0.09. The results indicated that it possessed the ~~kallikrein-protease~~ kallikrein production inhibition activity.

Please replace the paragraph beginning on page 9, line 23 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^6 virus/ml. A Chinese rabbit, weighing 1.5 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.1 ml per site in 250 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying with~~accompanied by changing colour from redness to mauveness, and the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra ~~dislocation and it was peeled in~~ after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 185 g, its SART activity was 0.71 iu/g, and its absorbance was 0.11. The results indicated that it possessed the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 10, line 14 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was

added ~~into~~ 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^7 virus/ml. A Blue-violet rabbit, weighing 2 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.2 ml per site in 100 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying with~~ accompanied by changing colour from redness to mauveness, ~~and~~ the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra ~~dislocation and it was peeled in~~ after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 235 g, its SART activity was 0.74 iu/g, and its absorbance was 0.13. The results indicated that it possessed the ~~kallikrein-protease~~ kallikrein production inhibition activity.

Please replace the paragraph beginning on page 11, line 5 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~ 500 ml PBS (-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^9 virus/ml. A Japanese white rabbit, weighing 3 kg, with its back shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.3 ml per

site in 200 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. ~~The injected rabbit injected-~~was fed for 3 days. ~~When the inflammatory tissue showed that the skin surface had visible blains aecompanying-withaccompanied by changing colour from redness to mauveness, and~~the skin became thick, and the subcuticle and the hip became swollen, the rabbit was killed by dislocating the cervical vertebra ~~disloeation-and it was peeled in-after~~ 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 335 g, its SART activity was 0.70 iu/g, and its absorbance was 0.12. The results indicated that it possessed the ~~kallikrein-protease~~kallikrein production inhibition activity.

Please replace the paragraph beginning on page 11, line 21 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~intoto~~ 500 ml of PBS (-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^6 virus/ml. A Japanese white rabbit, weighing 3 kg, with its back shaved, was sterilized with 75 % ethanol, injected subcutaneously with the virus injection of 0.1 ml per site in 200 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected-~~was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains aecompanying-withaccompanied by

changing colour from redness to mauveness, and the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra dislocation and it was peeled in after 15 min. The skin of the rabbit was packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 336 g, its SART activity was 0.61 iu/g, and its absorbance was 0.14. The results indicated that it possessed the ~~kallikrein-protease~~ kallikrein production inhibition activity.

Please replace the paragraph beginning on page 12, line 14 with the following:

The virus solution of the subculture antigen was taken from the -80°C refrigerator, and was thawed at 30°C in an incubator. Next, 5 ml of the virus solution was added ~~into~~ to 500 ml of PBS(-) by a 10-ml syringe, and was well shaken and diluted to an injection of 10^7 virus/ml. A Japanese white rabbit, weighing 3 kg, ~~with~~ while its back was shaved, was sterilized with 75% ethanol, injected subcutaneously with the virus injection of 0.2 ml per site in 200 sites with no leaking, no injecting without the virus injection and no puncturing throughout skin. The injected rabbit ~~injected~~ was fed for 3 days. When the inflammatory tissue showed that the skin surface had visible blains ~~accompanying~~ with accompanied by changing colour from redness to mauveness, and the skin became thick, and the subcuticle and the hip became swollen, ~~killed the rabbit~~ was killed by dislocating the cervical vertebra dislocation and it was peeled in after 15 min. The skin of the rabbit was

Serial No. 10/532,687

Office Action mailed March 5, 2007

Response filed April 5, 2007

packed in a plastic bag, and preserved at -18°C in a refrigerator prior to use. The skin of the rabbit weighed 335 g, its SART activity was 0.66 iu/g, and its absorbance was 0.12. The results indicated that it possessed the ~~kalikrein-protease~~ kallikrein production inhibition activity.